

WINSTROL DEPOT

COMPOSITION

Each ampoule contains:

Stanozolol 50 mg

Excipients:

- Sodium phosphate dibasic
- Polysorbate 80
- Sodium merthiolate
- Sodium chloride
- Water for injection

INDICATIONS

- General deterioration status, different origin slimness, anorexia not responding to treatment, convalescence, chronic and weakening diseases.
- Nephrotic and asthmatic syndromes, rheumatoid arthritis etc., for counteracting catabolic effects produced by corticosteroids.
- As coadjuvant in the treatment of decubitus ulcers, bone fractures of slow recovery, osteoporosis, extensive burns, periods before and after a surgical operation.
- In paediatrics, in growth failure in length and weight, somatic hypoevolutism, dystrophia and immaturity.

POSODOLOGY

Adults: 1 ampoule each 2 or 3 weeks.

Children:

- Up to 2 years: ½ ampoule each 2 or 3 weeks.
- From 2 to 6 years: ½ - 1 ampoule each 2 or 3 weeks, according to medical prescription.

Winstrol Depot is an aqueous suspension formulated for a prolonged absorption and with no local irritative effects. Its use is preferable provided that the doctor considers separate in time parenteral administrations more safe or more convenient than oral dairy administration.

Shake the ampoule before its use.

A diet rich and equilibrated is convenient when the product is being administered.

CONTRAINDICATIONS

Hypersensitivity to stanozolol,

Pregnancy

Prostate carcinoma

PRECAUTIONS

In sensitive patients to collateral endocrine or hidroelectrolitic (edema tendency) effects, a periodical vigilance in long term treatments is convenient.

In patients suffering hepatopathies, it is advisable to check cholestasis indexes. Winstrol tablets contain lactose. Intolerance to this ingredient has been reported in children and teenagers. Even the quantity present in this product is probably not sufficient to provoke the symptoms of intolerance, if diarrhoea appears you must consult your doctor.

WARNINGS

Anabolic steroids have the potential capacity of generating masculinizing effects, which can appear in the girls. If this happens, the treatment must be strictly controlled by the doctor. Long term treatment to promote growth must be interrupted when the skeletal age (to be controlled by radiology every 6 months) approaches the chronological age.

It is informed to sportsmen that this product contains a component that can give a positive analytical result in doping control.

INTERACTIONS

The simultaneous administration of corticosteroids or ACTH can occasionally provoke edema. In treatment with anticoagulants, an increase in the answer of these products can occur, and so it can be necessary to reduce the dosage to obtain the same therapeutic effect.

SIDE EFFECTS

They are rare and always reversible, consisting on nauseas, vomits, excitation, insomnia and acne

INTOXICATION AND TREATMENT

At the therapeutical dosages, no acute toxicity should be expected